



# Natural Antimicrobial Properties of *Hibiscus rosa-sinensis* Leaves and Their Potential in Eco-Friendly Agricultural Practices

Raman Jain\* and Aditya Panday<sup>2</sup>

<sup>1</sup> Department of Pharmacognosy, Bansal College of Pharmacy, Anand Nagar, Bhopal, Madhya Pradesh, India

<sup>2</sup> Akums Pharmaceutical Limited, Haridwar, Uttarakhand, India

Correspondence Author: Raman Jain

Received 24 May 2025; Accepted 5 Jun 2025; Published 19 Jun 2025

## Abstract

*Hibiscus rosa-sinensis* is a widely recognized plant known for its diverse phytochemical profile and traditional medicinal uses. The present study investigates the antioxidant and antibacterial potential of the ethanolic extract of *H. rosa-sinensis* leaves, which are deep green in color and rich in bioactive compounds. Antioxidant activity was evaluated using the DPPH free radical scavenging assay, where the leaf extract exhibited notable activity with an IC<sub>50</sub> value of 18.70 µg/mL, compared to the standard (ascorbic acid) IC<sub>50</sub> of 6.35 µg/mL. Antibacterial efficacy was assessed against both Gram-positive (*Staphylococcus aureus*) and Gram-negative (*Escherichia coli*) bacteria. The extract showed a concentration-dependent response, with higher antibacterial activity observed at 2% concentration compared to 1%. Notably, the extract demonstrated greater inhibition against *E. coli*, indicating stronger activity against Gram-negative strains. These findings suggest that *H. rosa-sinensis* leaves possess significant antioxidant and antibacterial properties and could serve as a promising source of natural bioactive agents for potential applications in sustainable agriculture, such as plant disease management and natural crop protection strategies.

**Keywords:** *Hibiscus rosa-sinensis*, antioxidant activity, antibacterial activity, ethanolic extract, sustainable agriculture

## Introduction

*Hibiscus rosa sinensis* vernacularly known as China rose or red Hibiscus is an average to large sized shrub. The plant supposed to native of Southeast Asia, belonging to mallow family Malvaceae. The plant is cultivated since long for their prominent beautiful flowers and medicinal attributes. Hibiscus flowers and leaves had been used in variety of condition like menorrhagia, emmenagogue, abortion, antifertility, contraceptive, demulcent, Cough. The leaves of the *Hibiscus* are ovate and alternatively arranged on the branches. The leaves would be intense green with serrate or toothed margins. The red flowered species would be preferred over others for therapeutic purpose.<sup>40</sup> the major constituents in the leaves are alkaloids, tannins, flavonoids, steroids, saponins, phenols, proanthocyanidin. The plant extract would be used in various skin disorders and protect the cells from damaging effect of oxidative stress and UV radiation. Due to presence of potential active metabolites, antioxidant and antibacterial activity of hydroalcoholic extract of leaves of *Hibiscus rosa sinensis* has been chosen for the study. Antioxidants are compounds which primarily neutralize the free radical in our body and protect the cell and tissue from invidious effects of oxidative stress.<sup>[1,2]</sup>

## Materials and methods

Analytical grade chemicals and reagents had been used for the purpose of study all the chemicals were procured from Central Drug House (P) LTD. New Delhi, the glassware used in the study was borosilicate and ASGI mark. Pharmaspec Shimadzu UV-VIS Spectrophotometer model UV-1700, Japan has been used.

## Collection and processing of plant material

The leaves of *Hibiscus rosa sinensis* has been collected from the herbal garden at BCP, Bhopal M.P. India, in the month of august. Plant sample were authenticated by Dr. S. Mishra, Scientist, MFP-PARK Laboratory, Bhopal, MP, India. The collected plant materials washed with tap water to remove debris and dirt. The plant was kept in shade until complete dried, the shade dried plant sample then grounded using electric grinder to get coarse powder, and the coarse powder was subjected to extraction with hydroalcoholic solvent.

## Extraction of plant material

The hydro alcoholic extract has been prepared by soaking the powdered sample in 80% ethanol. 250g of coarsely powdered plant sample was macerated with 80% ethanol for seven consecutive days in closed flask with periodically stirring. The extract was collected and filtered using Whatman filter paper, the filtrate was evaporated and concentrated to remove excess solvent under reduced pressure at 35°C in rotary evaporator. The concentrated extract was then placed in the desiccators to eradicate residual solvent.

## In-vitro anti-oxidant Activity

Antioxidant potential of *Hibiscus rosa sinensis* leaves extract had been evaluated by DPPH radical scavenging method. The comparison of sample data was made with ascorbic acid as standard antioxidant compound. 0.1mM solution of 2,2-Diphenyl-1-picrylhydrazyl (DPPH) was prepared in methanol<sup>[3]</sup>.

### Preparation of sample/standard

One mg of ascorbic acid and *Hibiscus rosa sinensis* dried powdered extract were dissolved in 1ml of methanol to get 1mg/ml standard and sample stock solution. Various dilution was made to get the desirable concentration of 20,40,60,80,100 µl/ml for both standard and sample in methanol. 2 ml of 0.1mM DPPH reagent was added and mixed thoroughly to each test tubes of sample and standard. The mixture is then incubated for 30 minutes in dark condition away from light then absorbance of standard and sample were recorded at the wavelength 517 nm <sup>[4, 5]</sup>.

### Preparation of control

Three milliliters of 0.1mM DPPH solution was prepared. The solution was incubated for 30 minutes at room temperature in dark condition. Absorbance of the control solution has been recorded against methanol as blank at 517 nm. The antioxidant activity of sample/ standard was reckoned by using formula <sup>[6, 7]</sup>.

Percentage Inhibition =  $\left[ \frac{\text{Abs of control} - \text{Abs of sample}}{\text{Abs of control}} \times 100 \right]$

### Antibacterial activity

Antimicrobial efficiency of the sample extract has been tested through well diffusion assay against gram negative bacteria *E. coli* MTCC 42 and gram-positive bacteria *S. aureus* MTCC 10787. The nutrient culture media was prepared by incorporation of twenty eight gram nutrient agar in one liter of distilled water. pH of media was checked after formulation the same has been recorded for future reference. The media was autoclaved at 121°C at 15 lbs pressure for 15 minutes, further sterilized media was allowed to cooled and poured into plates before it gets solidified all operation had been done in laminar flow <sup>[8-10]</sup>.

### Well diffusion assay

The sample for the test has been prepared by mixing 1% and 2% of test extract distinctly with distilled water. The culture of


bacterial strain was spread on prepared media. Standard solution for comparison with test has been prepared by dissolving one milligram of ofloxacin and gentamycin in 1ml of distilled water to get 1mg/1ml of standard solution. The inoculum of *E. coli* MTCC42 and *S. aureus* MTCC 10787 were prepared; primarily test organisms were inoculated in 10 mL of nutrient broth. The bacterial suspension was optimized to get 10<sup>8</sup> CFU/ml. 100 µl of the inoculum was taken and transferred in to pristine and sterile solidified agar media. Three wells of 6 mm had been made by sterile cork-borer. The initial two wells were filled with test sample with concentration of 1% and 2% further, third well were filled with 50µl of standard drug. The sample and standard were kept in sterile condition and allowed to diffuse for 30 minutes at room temperature further all samples were incubated for 24 hours at 37°C. The incubated plates were investigated for effect of test sample and standard. The clearing zone observed around the well was the harbinger of antimicrobial efficiency of tested compounds. The zone of inhibition was observed and measured in mm using ruler on the back of the inverted petri plate <sup>[11-14]</sup>.

**Table 1:** DPPH radical scavenging activity of ascorbic acid

Concentration (µg/ml)	Percentage inhibition
20	57.928
40	66.067
60	74.735
80	82.241
100	88.583
Control	0
IC50	6.35


**Table 2:** DPPH radical scavenging activity of *Hibiscus rosa sinensis*

Concentration (µg/ml)	Percentage inhibition
20	51.0570
40	62.790
60	68.498
80	77.589
100	79.809
control	0
IC50	18.70




**Fig 1:** Graph represents the percentage inhibition vs concentration of sample extracts**Table 3:** Antimicrobial activity of extract against *S. aureus*

Extract	Plate 1	Plate 2	Plate 3	Mean $\pm$ SD
1%	5 mm	6 mm	6mm	5.66 $\pm$ 0.533
2%	9 mm	10 mm	10 mm	9.66 $\pm$ 0.577
Control	0mm	0mm	0mm	0 $\pm$ 00
Ofloxacin (1mg/ml)	22 mm	20 mm	25 mm	22.33 $\pm$ 3.056

**Fig 2:** Antimicrobial activity of extract against *S. aureus***Table 4:** Antimicrobial activity of extract against *E. coli*

Extract	Plate 1	Plate 2	Plate 3	Mean $\pm$ SD
1%	11 mm	10 mm	8 mm	9.66 $\pm$ 1.231
2%	14 mm	13 mm	14 mm	13.33 $\pm$ 0.577
Control	0mm	0mm	0mm	0 $\pm$ 00
Gentamycin (1mg/ml)	25 mm	26 mm	24 mm	25 $\pm$ 1.487

**Fig 3:** Antimicrobial activity of extract against *E. coli*

## Result and discussion

The antioxidant activity of the ethanolic leaf extract was evaluated using the DPPH radical scavenging assay and compared with a standard. The extract demonstrated notable antioxidant capacity, with an IC<sub>50</sub> value of 18.70  $\mu$ g/mL, while the standard exhibited an IC<sub>50</sub> of 6.35  $\mu$ g/mL. Antibacterial efficacy was tested against *Staphylococcus aureus* (Gram-positive) and *Escherichia coli* (Gram-negative). The ethanolic extract showed a higher inhibitory effect against *S. aureus* at a 2% concentration, with a zone of inhibition measuring 9.66  $\pm$  0.577 mm, compared to 5.66  $\pm$  0.533 mm at 1%. Against *E. coli*, the extract was more effective at 2%, producing a zone of inhibition of 13.33  $\pm$  0.577 mm, as opposed to 9.66  $\pm$  1.231 mm at 1%. When comparing both bacterial strains, the extract displayed greater antibacterial activity against the Gram-negative *E. coli*. These results indicate that *Hibiscus rosa-sinensis* leaf extract possesses strong antioxidant and antibacterial properties, suggesting its potential use in bioactive formulations for similar applications.

## Conclusion

Increasing demand of herbal products inclined to explore the new potential substance from natural origin that could give better and safer alternative and efficiently used in various formulation. In this study we have tested the efficiency and potential of newer effects of *Hibiscus rosa sinensis* leaves extracts that might be useful to explore the plant in the newer area of herbal formulation. The results indicated that the *Hibiscus rosa sinensis* extracts had positive result on the parameter tested and could be used as anti-ageing, antibacterial and antioxidant potential. Further study needed to refine the extract by using isolated components and some more pharmacological evaluation needed that could broader the effect of drug.

## References

- Chin Hoong Fong. "Introduction, in The Hibiscus queen of tropical flowers Kuala Lumpur: Tropical Press, 1986, 2-3. Call no. RCLOS 635.93317 CHI.
- Sarjono Purbowatiningrum, Anggraeni Aulia, Monita Afiina, Asy'ari Mukhammad, Ismiyarto Ismiyarto, Ngadiwiyan Ngadiwiyan, etc. Nor. Antibacterial and Antioxidant Activity of Endophytic Bacteria Isolated from *Hibiscus tilaceus* Leaves. Jurnal Kimia Valensi. 2022;8:199-210. 10.15408/jkv.v8i2.25686.
- Al-Hashimi Alaa. Antioxidant and antibacterial activities of *Hibiscus sabdariffa* L. extracts. frican Journal of Food Science. 2012;6(21):506-511, 10.5897/AJFS12.099.
- Mak Yin, Chuah Li Oon, Rosma Ahmad, Bhat Rajeev. Antioxidant and antibacterial activities of hibiscus (*Hibiscus rosa-sinensis* L.) and Cassia (*Senna bicapsularis* L.) flower extracts. Journal of King Saud University - Science. 2013;25:275-282. 10.1016/j.jksus.2012.12.003.
- Wang J, Cao X, Jiang H, Qi Y, Chin KL, Yue Y. Antioxidant activity of leaf extracts from different *Hibiscus sabdariffa* accessions and simultaneous determination five major antioxidant compounds by LC-Q-TOF-MS. Molecules. 2014 Dec 17;19(12):21226-38. Doi: 10.3390/molecules191221226. PMID: 25525823; PMCID: PMC6271855.
- Hamrita B, Emira N, Papetti A, Badraoui R, Bouslama L, Ben Tekfa MI, *et al.* Phytochemical Analysis, Antioxidant, Antimicrobial, and Anti-Swarming Properties of *Hibiscus sabdariffa* L. Calyx Extracts: *In Vitro* and *In Silico* Modelling Approaches. Evid Based Complement Alternat Med. 2022 May 20;2022:1252672. Doi: 10.1155/2022/1252672. PMID: 35646135; PMCID: PMC9142284.
- Emesgen Assefa Abate, Alebel Nibret Belay. Assessment of antibacterial and antioxidant activity of aqueous crude flower, leaf, and bark extracts of Ethiopian *Hibiscus rosa-sinensis* Linn: geographical effects and Co<sub>2</sub>Res<sub>2</sub>/Glassy carbon electrode, International Journal of Food Properties. 2022;25(1):1875-1889. DOI: 10.1080/10942912.2022.2112598

8. Bukya Anil. A Comprehensive Analysis of Antioxidant Activity, Total Phenolic Content and Vitamin C From *Hibiscus Rosa Sinensis*. International journal of novel research and development. 2023;8(7):659-666.
9. Swargiary A, Daimari A, Daimari M, Basumatary N, Narzary E. Phytochemicals, antioxidant, and anthelmintic activity of selected traditional wild edible plants of lower Assam. Indian J Pharmacol. 2016 Jul-Aug;48(4):418-423. Doi: 10.4103/0253-7613.186212. PMID: 27756954; PMCID: PMC4980931.
10. Hamrita B, Emira N, Papetti A, Badraoui R, Bouslama L, Ben Tekfa MI, *et al*. Phytochemical Analysis, Antioxidant, Antimicrobial, and Anti-Swarming Properties of *Hibiscus sabdariffa* L. Calyx Extracts: *In Vitro* and *In Silico* Modelling.
11. Approaches. Evid Based Complement Alternat Med. 2022 May 20;2022:1252672. DOI: 10.1155/2022/1252672. PMID: 35646135; PMCID: PMC9142284.
12. Samsudin MS, Andriani Y, Sarjono PR, Syamsumir DF. “Study on Hibiscus Tiliaceus Leaves as Antibacterial and Antioxidant Agents”. Alotrop, 2019 Dec, 3(2). DOI:10.33369/atp.v3i2.9874.
13. Ruban P, Gajalakshmi K. In vitro antibacterial activity of Hibiscus rosa-sinensis flower extract against human pathogens. Asian Pac J Trop Biomed. 2012 May;2(5):399-403. DOI: 10.1016/S2221-1691(12)60064-1. PMID: 23569938; PMCID: PMC3609315.
14. Anna Ricka J Barraquia, Ann Patrice L Cabuso, Charmaine M. Narvacan. Antibacterial Activity of Ethanolic Extract of Hibiscus Rosa-Sinensis Flower Against Staphylococcus Epidermidis and Staphylococcus Saprophyticus, Lyceum of the Philippines-St. Cabrini College of Allied Medicine Research. 2017 March;2(2):39-50.